



**WE CARE FOR you,
AND YOUR NEW ARRIVAL**

In infertility associated with PCOS

LETZY

Letrozole Tablets USP 2.5 / 5 mg

SAFER

BETTER

ECONOMICAL




Inhibit

Depletion of
ER receptor



High

Pregnancy
Rate



Ensures

Monofollicular
Rate

Dosage :

Anovulatory Infertility,
PCOS - 2.5 mg OD from day 3 to day 7



In Emesis,

MONIDOX-XT

Doxylamine Succinate 20 mg (10 mg IR & 10 mg ER)
+ Pyridoxine Hydrochloride 20 mg (10 mg IR & 10 mg ER)
+ L-Methylfolate Calcium 1 mg & Methylcobalamin 1500 mcg.

BE SAFE WITH THE NO. 1 ANTIEMETIC

10_{mg}

Immediate Release

Rapid Onset of
Action

10_{mg}

Enteric Coated Core

Long Lasting
Therapeutic Effects



Benefits for Mother

Prevents Emesis &
Progression to HEG*



Foetal Safety^{1,2}

No Risk of Cleft Palate,
Cleft Lip & Septal Defects

Dosage :

1 Tab at night or 1 Tab in morning



Control NVP & Prevent NTDs,

MONIDOX PLUS

Doxylamine Succinate 20 mg + Pyridoxine Hydrochloride 20 mg + Folic Acid 5 mg.

The No. 1 Anti-emetic Fortified with Folic acid




Ondansetron

Treatment During
Pregnancy may cause
birth defects



Benefits for Mother

Adequate Folic Acid
supplementation
during pregnancy



Doxylamine Succinate

Blocks Chemoreceptor
trigger zone & controls
Nausea & Vomiting




Foetal Safety

Prevent NTDs



Pyridoxine

Restores Tryptophan
Metabolism



In Iron Deficiency Anaemia During Pregnancy & Lactation

HBMER-XT

Ferrous Ascorbate equivalent to elemental Iron 100 mg, Folic Acid 1.5 mg & Elemental Zinc equivalent to 22.5 mg Tablets

WHO recommends ZINC supplement during pregnancy

Zn

Zinc

Reduces IUGR and
boost the Immunity

B₉

Folic Acid

Helps in
RBC Formulation

FA

Ferrous Ascorbate

Provides faster rise
in hemoglobin



In Iron Deficiency Anaemia,
During Pregnancy & Lactation

HBMER-Z

Ferrous Fumarate 300 mg + Folic Acid 750 mcg + Zinc Sul. 61.8 mg.

The Joy of a perfect Motherhood

Zn

Zinc

Improves
Immunity of
Mother & Child

B₉


Folic Acid

Prevents
Neural Tube
Defect

B₁₂

Vitamin B12

Improves
Pregnancy
Outcome



In Spasm associated with Pain

MVERIN


Drotaverin 80 mg + Mefenamic Acid 250 mg. Tablets

Distinct Spasmolytic Analgesic Solution



Drotaverine

Relief from
abdominal pain



Mefenamic Acid

Powerful anti
inflammatory
& Analgesic



In Dysmenorrhea,

DISPAS

Mefenamic Acid IP 250 mg. + Dicyclomine Hydrochloride IP 10 mg.

Fast Action Against Dysmenorrhea



Mefenamic Acid

Reduces pain



Dicyclomine

Offers relief from
cramps




In Male Infertility,
Decreased sperm count & SUBOPTIMAL
sperm parameter are responsible



SUPERMAX+

Clomifene Citrate 25 mg. & Antioxidants, Multivitamins, Amino Acids & Minerals

Complete Fertility Enhancer




CC-25 mg
Co-Q-10 to 60 mg

Influences Spermatogenesis
Increase sperm count
& sperm motility



Lycopene - 4 mg
Levocartinine - 50 mg

Anti Oxidants & Amino Acid
Improve Sperm Motility



B6 - 1.5 mg, B9 - 5 mg
B12 - 1500 mcg

Multi Vitamins
Improve sperm quality



Zinc - 66 mg
Minerals

Essential for Testosterone
production &
Sperm morphology



Selenium - 200 mcg
Minerals

Essential for Testosterone
production

Dosage :

25 days - White Tablets &
30 days - Brown Tablets
3 months therapy for maximum result

Supermax+ Tablets

Indication

- Male Infertility – To improve Sperm count, Sperm Motility & Sperm Morphology

Dosage

- 1 WHITE Tablet of Clomiphene Citrate for 25 days with gap of 5 days &
- 1 BROWN Tablet of Antioxidant for 30 days
- For 3 months or as directed by the Physician

Because Spermatogenesis takes 3 months

Mode of Action -- Simplified

Increases Sperms Concentration	Increases Sperms Morphology	Increases Sperms Motility
Clomiphene Citrate		
Coenzyme Q10		Coenzyme Q10
Lycopene	Lycopene	Lycopene
		Levocarnitine
Vitamin B6, B9, B12		
Zinc, Selenium	Zinc	

Free Radical Fighting Foods for Male infertility

All of the cells of the human body are susceptible to free radical damage, including sperm cells. Studies have shown that 40 percent of infertile men have high levels of free radicals, Vitamins C and E, and the mineral Selenium provide strong antioxidant properties to fight free radical damage and maintain the integrity of sperm cells. To bolster your system and combat injury from free radicals, choose foods from the following :

Foods High in Vitamin C :

All fruits and vegetables, particularly Cherry, Broccoli, Citrus fruits, Black currants, Green leafy vegetables, Guavas, Kiwi, Mangoes, Melons, honeydew, watermelon, papayas, peppers, strawberries, Tomatoes.

Foods High in Vitamin E

Almonds, Blackberries, Cod liver oil, Corn, corn oil, Cottonseed oil, Currants (Black) Grapefruit, Oats, Olive oil, Peanuts, Safflower oil, Sunflower seeds/Sunflower oil, wheat and wheat germ.

Foods High in Selenium

Onions Bran, Spices, Brazil nuts, Tomatoes Eggs, Turnips, Fish & seafood, Vinegar, Wheat germ, Whole grains, Mushrooms.



**Each brown colour tablet contains :
Antioxidants, Multivitamins, Amino Acid
& Minerals Tablets**

Ubidecarenone USP	60 mg
Zinc Suphate IP	66 mg
Folic Acid IP	5 mg
Methylcobalamin JP	1500 mcg
Pyridoxine Hydrochloride IP	1.5 mg
Lycopene 6% USP	4 mg
Selenium	
(As Sodium Selenite BP)	200 mgc
Levocarnitine L-Tartrate	50 mg




**Each white colour tablet contains :
Clomifene Citrate Tablet IP
Clomifene Citrate IP 25 mg**

In Inflammation, Injury, Wound, Osteoarthritis and Cellulitis

BRUZYME


Trypsin 48 mg. + Bromelain 90 mg. + Rutoside Trihydrate 100 mg.

Trusted Healer, Internally & Externally




Trypsin

Better Resolution
of inflammatory
Symptoms



Bromelain

Reducing pain
and Swelling
Especially of the
nose and sinuses



Rutoside

Help strengthen
and increase flexibility
in blood vessels

Dosage :

2 Tab 3 Times a Day
30 min. before meal
or 2 hrs. after meal




For Instant Control

MEPAN-DSR


Pantoprazol 40 mg + Domperidon Prolonged Release Capsule 30 mg

Well Suited for elderly patients & Cardiac patients on multiple therapy



Pantoprazol


Reduce the
amount of acid
your stomach makes



Domperidon

increase the
movements or contractions
of the stomach and bowel

Dosage :
1 capsule
before breakfast



For, Asthmatic Cough

SYP SAL-AG

Levosalbutamol 1 mg. + Guaiphenesin 50 mg.
+ Ambroxal 30 mg. + Menthol Syrup

The Right Choice in Your Asthmatic Cough Patients




Levosalbutamol

is used to treat
Asthma & Chronic
obstructive pulmonary
disease (COPD)



Ambroxol

is a secretolytic agent
used in the treatment
of respiratory disease




Guaiphenesin

increase the
efficiency of the
cough reflex and
facilitate removal
of the secretions



Dosage :
2 Table Spoon
After Lunch & Dinner



In, Infection of the Respiratory Tract, Ear, Nose, Throat

TAB-MUCILLIN-375

Sultamicillin 375 mg.

In Obstetric and gynaecological infection and in the treatment of gonorrhoea

Mucillin-375 has been shown to be clinically effective in non-comparative trials in patients with infections of the respiratory tract, ears, nose and throat, urinary tract, skin and soft tissues, as well as in obstetric and gynaecological infections, and in the treatment of gonorrhoea.

Mode of Action of Sultamicillin :

Sultamicillin interrupts bacterial cell wall synthesis, which is the protective covering of bacteria. This leads to death of bacterial cells, thereby offering effective infection control.

Dosage :

1 Tab at night or 1 Tab in morning

